

MICROPIGMENTATION PRE PROCEDURE ADVICE

Please read the following advice carefully and sign at the end

- Microblading procedure normally requires multiple treatment sessions. For best results, clients will be required to return for at least one re-touch appointment. This will take place 5- 6 weeks after the initial procedure. Those with oily skin may require an additional touch up. Please be aware that color intensity will be significantly darker and sharper immediately and a few days after the initial procedure, but the color will reduce by 30-50%
- Although numbing cream is used during the procedure, sensitivity or discomfort may still be felt. Skin may be red and/or swollen after the procedure
- Please do not drink alcohol 24 hours prior to the treatment
- A patch test will be performed, unless waived by client
- Please do not shape or wax your brows before the procedure. Your technician will shape brows during the procedure
- No electrolysis for at least 5 days before the procedure
- Botox, AHA products and retinoids should be avoided for 2 weeks prior to the procedure
- Exfoliating treatments such as microdermabrasion should not be performed within 2 weeks prior to procedure
- Chemical and laser peels should be avoided no less than 6 weeks prior to procedure
- Patients prone to cold sores/fever blisters should take an anti-viral prior to treatment
- Hormone therapies can affect pigmentation and/or cause sensitivity

Topical Anaesthetic Advice

- **Allergic reaction** can occur from any anaesthetics used during the procedure. If you do suffer from an allergic reaction, you should contact your doctor immediately. Allergic reaction response may show through redness, swelling, rash, blistering, dryness or any other symptoms associated with an allergic reaction.
- **Numbness** – We cannot accept responsibility if the area to be treated does not respond to the numbing cream. Each individual is different according to skin type. Some clients report the area to be completely numb, while others may experience some discomfort.
- **Procedure** – For microblading procedure, a numbing cream/gel is used. The products are formulated to be perfectly safe and can be purchased over the counter from any pharmacy/chemist. The anaesthetic is placed over the treatment area for 20-30 minutes then carefully removed prior to treatment. As a result of the treatment, combined with the use of the anaesthetic, you can expect to experience some redness/swelling that can last 1-4 days. You should always follow your post procedure advice and after care for the best results.

Contraindications for Microblading

- Liver disease – high risk of infection
 - Compromised skin near brow area
 - Pregnancy/Nursing
 - Chemotherapy/Radiation
- **The following medical conditions require a note from your doctor giving consent**
Diabetes Type 1 and 2, high blood pressure, auto-immune disease, thyroid / Graves' disease Any other medical condition that causes slow healing or a high risk of infection

I have read and full understood the above information provided and any risks involved with the use of topical anesthetic and I therefore consent to the use of the anesthetic for the microblading procedure. I agree to follow pre- and post-procedure advice closely

Client name: _____ Signature: _____ Date: _____